

National Academies Press © 2013

U.S. HEALTH IN INTERNATIONAL PERSPECTIVE:
SHORTER LIVES, POORER HEALTH

Grantmakers in Health April 10, 2013




Motivations


- Prior NRC (2011) report "Explaining Divergent Levels of Mortality in High-Income Countries" (sponsored by NIA)
- Current NRC/IOM (2013) report looking at morbidity and mortality at all ages (sponsored by OBSSR and others)
- Value of cross-national perspective for understanding US health



Overview of Comparison Group




- 17 comparable highincome nations
- Australia, Austria,
 Canada, Denmark,
 Finland, France, Germany,
 Italy, Japan, Norway,
 Portugal, Spain, Sweden,
 Switzerland, the
 Netherlands, the United
 Kingdom, and the United
 States




The Health of the Nation: Poor Value for Money

Life Expectancy and Health Care Spending in OECD Countries (2008)



Or latest year available.
 Source: OECD Health Data 2010.








Mortality Rates in Peer Countries, 2008


Non-communicable diseases

Communicable diseases









Higher Mortality Rates and Lower Life Expectancy

Mortality Rates by Cause of Death

	Mortali	ty Rate (per 100,000)	
		Peer Countries (N = 16)
Cause of Death	United States	Unweighted Mean	Range
	U.S. Death Rates		
	Above Average		
Cardiovascular diseases ^a	155.7	133.6	97.4-174.9
Neuropsychiatric conditions ^b	39.2	28.1	7.2-48.4
Respiratory disease	34.3	21.0	12.7-34.4
Infectious and parasitic diseases	15.4	7.7	4.4-17.5
Diabetes mellitus	15.2	10.2	4.5-19.3
Genitourinary diseases ^c	12.3	7.2	3.0-12.2
Endocrine disorders	7.1	4.2	1.6-8.1
Congenital anomalies	4.3	3.3	2.6-4.0
Musculoskeletal diseases ^d	2.9	2.4	1.2-3.5
Nutritional deficiencies	1.0	0.7	0.1-2.0
Skin diseases	0.8	0.6	0.1-1.5
Maternal conditions	0.4	0.1	0.0-0.2
Perinatal conditions	7.1	3.7	1.3-5.9
Unintentional injuries ^c	35.5	20.4	13.7-38.6
Intentional injuries	17.3	11.4	5.6-20.2
	U.S. Death Rates at or Below Average		
Malignant neoplasms ^f	123.8	127.3	106.5-157.7
Digestive diseasesg	19.8	19.8	13.0-29.5
Respiratory infections	9.7	12.3	4.0-29.7
Other neoplasms	2.9	3.3	2.1-5.3
Oral conditions ^h	0.0	0.0	0.0-0.1
Sense organ diseases ^h	0.0	0.0	0.0

Life Expectancy


Ma	ales		Fen	nales	
Country	LE	Rank	Country	LE	Rank
Switzerland	79.33	1	Japan	85.98	1
Australia	79.27	2	France	84.43	2
Japan	79.20	3	Switzerland	84.09	3
Sweden	78.92	4	Italy	84.09	3
Italy	78.82	5	Spain	84.03	5
Canada	78.35	6	Australia	83.78	6
Norway	78.25	7	Canada	82.95	7
Netherlands	78.01	8	Sweden	82.95	7
Spain	77.62	9	Austria	82.86	9
United Kingdom	77.43	10	Finland	82.86	9
France	77.41	11	Norway	82.68	11
Austria	77.33	12	Germany	82.44	12
Germany	77.11	13	Netherlands	82.31	13
Denmark	76.13	14	Portugal	82.19	14
Portugal	75.87	15	United Kingdom	81.68	15
Finland	75.86	16	United States	80.78	16
United States	75.64	17	Denmark	80.53	17






Changes in Life Expectancy over Time

Life Expectancy at Birth - Male





Life Expectancy at Birth - Female




Probability of Survival to Age 50 - Male

Probability of Survival to Age 50 - Female








THE NATIONAL ACADEMIES


Advisers to the Nation on Science, Engineering, and Medicine

Low Life Expectancy at Every Age

Ranking of US Mortality Rates by Age Group vs. Peer Countries, 2006-2008




For either sex, the US rank is never better than 15th out of 17 until age 75.






Low Life Expectancy for Non-Hispanic Whites

Ranking of Mortality of Non-Hispanic Whites by Age Group vs. Peer Countries, 2006-2008




For either sex, the US rank is never ranked higher than 16th out of 17 until age 55.







BMI and Obesity

Average BMI in Peer Countries, 2008





- The US has the highest prevalence of adult obesity among all OECD countries, a position it has held for decades.
- As of 2009, the prevalence of obesity in the US was twice the OECD average.






Diabetes

Self-Reported Diabetes Prevalence in Peer Countries, 2008






As of 2010, the US had close to the highest prevalence of diabetes in adults aged 20-79 in all peer countries (and all OECD countries except Mexico).






Infant Mortality in Peer Countries

Deaths per 1,000 Live Births






- From 2005-2009, the US had the highest infant mortality rate of the peer countries and the 9th highest in the OECD.
- Well-educated non-Hispanic whites and mothers have higher infant mortality rates in the US than those in other countries.
- Since the 1970s, US infant mortality has not kept pace with declines achieved by other countries.
 - US infant mortality declined by 20% from 1990-2010, but comparable high-income countries halved their rates.




Other Birth Outcomes

Low Birth Weight in Peer Countries



Global Preterm Birth Rates



- The US prevalence of low birth weight is the second highest of the peer countries.
- US stillbirths and perinatal mortality rates are also among the highest.
- A 2012 analysis found that the US preterm birth rate (12%) was comparable to that of sub-Saharan Africa.



Blencowe et al. 2012

THE NATIONAL ACADEMIES

Advisers to the Nation on Science, Engineering, and Medicine

US Children's Health Disadvantage


Children in the US have the **highest probability of dying before**age 5 of any of the peer countries.



- In 2004, 11% of US deaths before age 5 were from injuries.
- In 2006, the US had the highest rate of child deaths due to negligence, maltreatment, or physical assault.
- The violent death rate among US boys aged 1-4 has exceeded the OECD average since the late 1960s
- The US is ranked 24th of 30 (OECD) and 21 of 21 (UNICEF) on selected measures of children's well-being.

US Adolescent Health Disadvantage

Prevalence of overweight (including obese) children in peer countries (2011)





- Adolescent obesity rates in the US far exceed those in comparable nations
- In 2010, the US had the 5th highest prevalence of diabetes among children ages 0-14 in peer countries.
- Among teens aged 15-19 in 2005, the US had the highest all-cause mortality rate among peer countries.

Adolescent Sexual Health

- The US has the highest teen pregnancy rate, nearly 3.5 times the average of peer countries in 2010.
- The US has a higher prevalence of syphilis, gonorrhea, and chlamydia among 15-19 year-olds than any other high-income country that provided comparison data.
- Among high-income countries, the
 US has the highest prevalence of
 HIV infection at ages 15-24.

Adolescent Birth Rate, 2010










Youth Injury Mortality

Transportation-related Mortality in Peer Countries






- Among adolescents aged 15-19 in 2005, the US had the highest injury mortality rate.
- Since the 1950s, transport injury mortality at ages 15-24 has been higher in the US than in peer countries.
- Since the 1960s, the US has had higher non-transport injury mortality among children aged 5-9, and especially among males aged 10-19.






Youth Mortality from Violence

Violent Mortality in Peer Countries





- Since the 1950s, males aged 15-24 have been far more likely to die from violence in the US than in the peer group.
- US males aged 15-19 are 5 times more likely to die from violence than those in other OECD countries.





Age 50 and Above

Disease and Disadvantage in the United States and in England

Michael M	farmot, MI
Zoe Oldfie	AL MSc

Contest: The United States spends considerably more money on health care than the United Kingdom, but whether that translates to better health outcomes is unknown. Objective: To asserts the reliable health status of older individuals in fingular and the United States, repossibly how their health status varies by important indication of sooloeconomic position.

Reales Catting and Particles are We undered reconstricts commiss of rest.

Differences In Disease Prevalence As A Source Of The U.S.-European Health Care Spending Gap A Comparison of the He With That of Europeans | Maurico Averdano, Proj. M. Maria Glymoux, Sc. A Tale of Two Countries—the Unit

Americans are diagnosed with and treated for several chronic illnesses more often than their European counterparts are.

by Kenneth E. Thorpe, David H. Howard, and Katya Galactionova

EXPLAINING DIVERGENT LEVELS OF LONGEVITY IN HIGH-INCOME COUNTRIES

Eileen M. Crimmins, Samuel H. Preston, and Barney Cohen Editors

Panel on Understanding Divergent Trends in Longevity in High-Income Countries

Health Disadvantage in US Adults Aged 50 to 74 Years: A Comparison of the Health of Rich and Poor Americans With That of Europeans

Mauricio Avendano, PhD, M. Maria Glymour, ScD, James Banks, PhD, and Johan P. Mackenbach, PhD

A Tale of Two Countries—the United States and Japan: Are Differences in Health Due to Differences in Overweight?

Sandra L. Reynolds¹, Aaron Hagedorn², Jihye Yeom², Yasuhiko Saito³, Eise Yokoyama³, and Eileen M. Crimmins²

¹University of South Florida, Tampa, FL, USA


²University of Southern California, Los Angeles, CA, USA

³Nihon University, Tokyo, Japan

Differences in health between Americans and Western Europeans: Effects on longevity and public finance

Pierre-Carl Michaud a.*, Dana Goldman b, Darius Lakdawalla b, Adam Gailey c, Yuhui Zheng d

- Conditions more prevalent in the US among those age 50 and older
 - Obesity
 - Diabetes
 - Hypertension
 - Heart disease
 - Myocardial infarction
 - Stroke
 - Chronic lung disease
 - Asthma
 - Cancer
 - Arthritis
 - Activity limitations



THE NATIONAL ACADEMIES

Advisers to the Nation on Science, Engineering, and Medicine

Nine Areas of US Health Disadvantage

Infant Mortality & Low Birth Weight

Injuries & Homicides

Adolescent
Pregnancy &
STIs


HIV & AIDS

Drug-related
Deaths


Obesity & Diabetes


Cardiovascular Disease Chronic Lung
Disease

Disability



Structure of the Report





Determinants of Health

Health Systems

- Insurance and access barriers
- Weak primary care
- Coordination and errors

Individual Behaviors


- Caloric intake
- Drug use
- Sexual practices
- Driver safety
- Firearm possession

The **Environment**


- Physical environment
- Social environment

Social Factors

- □ Low income/poverty
- Education
- Social mobility
- Incarceration
- Single-parent households




Policies and Social Values



Switzerland United States Australia Japan Canada Spain Portugal Netherlands United Kingdom Norway Italy Finland Germany Austria France Denmark Sweden 15 0 20 30 35 40 Percentage of Gross Domestic Product

Bradley E H et al. BMJ Qual Saf 2011;20:826-831




THE NATIONAL ACADEMIES

Advisers to the Nation on Science, Engineering, and Medicine


Research Recommendations

- Recommendation 1: NIH and NCHS should join with an international partner to improve the quality and consistency of data sources available for cross-national health comparisons.
- Recommendation 2: NIH and other research funding agencies should support the development of more refined methods and study designs for cross-national health research.
- Recommendation 3: NIH and other research funding agencies should commit to a coordinated portfolio of investigator-initiated and invited research devoted to understanding the factors responsible for the U.S. health disadvantage and potential solutions, including lessons that can be learned from other countries.




Policy Recommendations

- Recommendation 4: Intensify efforts to achieve established national health objectives that are directed at the specific disadvantages documented in this report and that use strategies and approaches that reputable review bodies have identified as effective.
- Recommendation 6: Commission a review of the available evidence on (1) the effects of policies on the areas in which the US has an established health disadvantage, (2) how these policies have varied over time across high-income countries, and (3) the extent to which these policy differences may explain cross-national health differences.



Recommendation for Philanthropy

Recommendation 5: The philanthropy and advocacy communities should organize a comprehensive media and outreach campaign to inform the general public about the U.S. health disadvantage and to stimulate a national discussion about its implications for the nation.



Next Steps

- Disseminate widely
- Stimulate a national conversation
- Conduct additional research
- Support sound policies



Contact Information

- Steven H. Woolf, MD, MPH
 Director, VCU Center on Human Needs
 Virginia Commonwealth University
 swoolf@vcu.edu
- Laudan (Laudy) Aron, MA
 Senior Research Associate, Urban Institute
 202-261-5369, LAron@urban.org

